

兔抗 PPBP 多克隆抗体

中文名称: 兔抗 PPBP 多克隆抗体

英文名称: Anti-PPBP rabbit polyclonal antibody

别名: PBP; TC1; TC2; TGB; LDGF; MDGF; TGB1; B-TG1; CTAP3; CXCL7; NAP-2; SCYB7; THBGB; LA-PF4; THBGB1; Beta-TG; CTAPIII; CTAP-III

储存: 冷冻 (-20℃)

抗原: PPBP

宿主: Rabbit

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

相关类别: 一抗

技术规格

Background:	The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E ₂ secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. The protein also is an antimicrobial protein with bactericidal and antifungal activity.
Applications:	ELISA, WB, IHC

Name of antibody:	PPBP
Immunogen:	Synthetic peptide of human PPBP
Full name:	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7)
Synonyms :	PBP; TC1; TC2; TGB; LDGF; MDGF; TGB1; B-TG1; CTAP3; CXCL7; NAP-2; SCYB7; THBGB; LA-PF4; THBGB1; Beta-TG; CTAPIII; CTAP-III
SwissProt:	P02775
ELISA Recommended dilution:	2000-10000
IHC positive control:	Human thyroid cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	14 kDa
WB Positive control:	Human plasma solution
WB Recommended dilution:	1000-5000



