

兔抗 SEPT7 多克隆抗体

中文名称: 兔抗 SEPT7 多克隆抗体

英文名称: Anti-SEPT7 rabbit polyclonal antibody

别 名: CDC3; CDC10; SEPT7A; NBLA02942

相关类别: 一抗

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: SEPT7

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

This gene encodes a protein that is highly similar to the CDC10 protein of S accharomyces cerevisiae. The protein al so shares similarity with Diff 6 of Dros ophila and with H5 of mouse. Each of these similar proteins, including the ye



tif. The yeast CDC10 protein is a struct ural component of the 10 nm filament which lies inside the cytoplasmic mem brane and is essential for cytokinesis. This human protein functions in gliom agenesis and in the suppression of gli oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19. Applications:
which lies inside the cytoplasmic mem brane and is essential for cytokinesis. This human protein functions in gliom agenesis and in the suppression of gli oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
brane and is essential for cytokinesis. This human protein functions in gliom agenesis and in the suppression of gli oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
This human protein functions in gliom agenesis and in the suppression of gli oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
agenesis and in the suppression of gli oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
oma cell growth, and it is required for the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
the association of centromere-associate d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
d protein E with the kinetochore. Alter native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
native splicing results in multiple trans cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
cript variants. Several related pseudoge nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
nes have been identified on chromoso mes 5, 7, 9, 10, 11, 14, 17 and 19.
mes 5, 7, 9, 10, 11, 14, 17 and 19.
Applications: WB
Name of antibody: SEPT7
Synthesized peptide derived from inter Immunogen:
nal of human SEPT7.
Full name: septin 7
Synonyms: CDC3; CDC10; SEPT7A; NBLA02942
SwissProt: Q16181



www.shjning.com

WB Predicted band size:	51 kDa
WB Positive control:	MCF-7 cell lysate
WB Recommended dilution:	500-3000

