

## 兔抗 RPL17 多克隆抗体

- 中文名称：兔抗 RPL17 多克隆抗体
- 英文名称：Anti-RPL17 rabbit polyclonal antibody
- 别名：ribosomal protein L17; L17; PD-1; RPL23
- 相关类别：一抗
- 储存：冷冻（-20℃）
- 宿主：Rabbit
- 抗原：RPL17
- 反应种属：Human, Mouse, Rat
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

### 技术规格

**Background:**

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22P family of ribosomal proteins. It is located in the cytoplasm. This gene has been referred to as rpL23 because the encoded protein shares amino acid identity with ribosomal protein L23 from *Halobacterium marismortui*; however, its official symbol is RPL17. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Alt

	ernative splicing results in multiple transcript variants. Read-through transcription also exists between this gene and the neighboring downstream C18orf32 (chromosome 18 open reading frame 32) gene.
<b>Applications:</b>	ELISA, WB, IHC
<b>Name of antibody:</b>	RPL17
<b>Immunogen:</b>	Fusion protein of human RPL17
<b>Full name:</b>	ribosomal protein L17
<b>Synonyms:</b>	L17; PD-1; RPL23
<b>SwissProt:</b>	P18621
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human tonsil
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	21 kDa
<b>WB Positive control:</b>	293T cell, Mouse brain tissue, Mouse testis tissue, HepG2 cell, Mouse liver tissue, A375 cell lysates
<b>WB Recommended dilution:</b>	1000-5000



