

## 兔抗 SIGLEC12 多克隆抗体

中文名称: 兔抗 SIGLEC12 多克隆抗体

英文名称: Anti-SIGLEC12 rabbit ployclonal antibody

别 名: S2V; SLG; SIGLECL1; Siglec-XII

相关类别: 一抗

储 存: 冷冻(-20℃)

宿 主: Rabbit

抗原: SIGLEC12

反应种属: Human, Mouse

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

Background:	Sialic acid-binding immunoglobulin-like lectins (SIG LECs) are a family of cell surface proteins belongin g to the immunoglobulin superfamily. They mediat e protein-carbohydrate interactions by selectively b inding to different sialic acid moieties present on glycolipids and glycoproteins. This gene encodes a member of the SIGLEC3-like subfamily of SIGLECs. Members of this subfamily are characterized by an extracellular V-set immunoglobulin-like domain foll owed by two C2-set immunoglobulin-like domains, and the cytoplasmic tyrosine-based motifs ITIM an d SLAM-like.
Applications:	ELISA, WB, IHC
Name of antibody:	SIGLEC12
Immunogen:	Fusion protein of human SIGLEC12



## www.shjning.com

Full name:	sialic acid binding Ig-like lectin 12 (gene/pseudoge ne)
Synonyms:	S2V; SLG; SIGLECL1; Siglec-XII
SwissProt:	Q96PQ1
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human prostate cancer
IHC Recommend dilution:	25-100
WB Predicted band size:	65 kDa
WB Positive control:	Mouse bladder tissue
WB Recommended dilution:	200-1000





