

兔抗 STK40 多克隆抗体

中文名称: 兔抗 STK40 多克隆抗体

英文名称: Anti-STK40 rabbit polyclonal antibody

别 名: SHIK; SgK495

相关类别: 一抗

抗 原: STK40

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	STK40 (Serine/threonine-protein kinase 40), also known as Sugen kinase 495, is a 435 amino acid protein kinas e that may be a negative regulator of NFκB and p53-me ed gene expression. Localized to both the cytoplasm and the nucleus, STK40 is ubiquitously expressed. STK40 has a central serine/threonine protein kinase domain and is homologous to TRB-3, a protein that regulates activation of MAP kinases and inhibits NFκB-mediated gene transcription.
Applications:	ELISA, WB, IHC
Name of antibody:	STK40
Immunogen:	Fusion protein of human STK40
Full name:	serine/threonine kinase 40
Synonyms :	SHIK; SgK495



SwissProt:	Q8N2I9
ELISA Recommended dilution:	1000-2000
IHC positive control:	Human cervical cancer and human thyroid cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	49 kDa
WB Positive control:	Human placenta tissue
WB Recommended dilution:	200-1000





