

兔抗 SPIN1 多克隆抗体

- 中文名称: 兔抗 SPIN1 多克隆抗体
- 英文名称: Anti-SPIN1 rabbit polyclonal antibody
- 别 名: SPIN
- 抗原: SPIN1
- 储存: 冷冻(-20℃) 避光
- 宿 主: Rabbit
- 反应种属: Human Mouse Rat
- 相关类别: 一抗
- 标记物: Unconjugate
- 克隆类型: Unconjugate

技术规格

Background:	Chromatin reader that specifically recognizes and bin ds histone H3 both trimethylated at 'Lys-4' and asy mmetrically dimethylated at 'Arg-8' (H3K4me3 and H 3R8me2a) and acts as an activator of Wnt signaling pathway dowstream of PRMT2. In case of cancer, pr omotes cell cancer proliferation via activation of the Wnt signaling pathway (). Overexpression induces m etaphase arrest and chromosomal instability. Localize s to active rDNA loci and promotes the expression of rRNA genes (). May play a role in cell-cycle regul ation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process th at takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulatin g maternal transcripts to control meiotic resumption.
-------------	---



www.shjning.com

Applications:	WB
Name of antibody:	SPIN1
Immunogen:	Synthesized peptide derived from internal of human SPIN1.
Full name:	spindlin 1
Synonyms :	SPIN
SwissProt:	Q9Y657
WB Predicted band size:	30 kDa
WB Positive control:	Hela cells lysate
WB Recommended dilution:	500-3000





