

## 兔抗 STARD10 多克隆抗体

中文名称: 兔抗 STARD10 多克隆抗体

英文名称: Anti-STARD10 rabbit polyclonal antibody

别 名: PCTP2; CGI-52; NY-CO-28; SDCCAG28

相关类别: 一抗

存: 冷冻(-20℃) 避光

宿 主: Rabbit

抗 原: STARD10

反应种属: **Human Mouse Rat** 

标 记 物: Unconjugate

克隆类型: Unconjugate

## 技术规格

Background:	May play metabolic roles in sperm maturation or ferti lization By similarity. Phospholipid transfer protein that t preferentially selects lipid species containing a palmitoyl or stearoyl chain on the sn-1 and an unsaturated fatty acyl chain (18:1 or 18:2) on the sn-2 position. A ble to transfer phosphatidylcholine (PC) and phosphatidylcholamline (PE) between membranes.
Applications:	WB
Name of antibody:	STARD10
Immunogen:	Synthesized peptide derived from N-terminal of huma n STARD10.
Full name:	StAR-related lipid transfer (START) domain containing 10
Synonyms:	PCTP2; CGI-52; NY-CO-28; SDCCAG28



## www.shjning.com

SwissProt:	Q9Y365
WB Predicted band size:	33 kDa
WB Positive control:	COLO205 cells lysate
WB Recommended dilution:	500-3000

