

FCGR3A 抗原（重组蛋白）

中文名称： FCGR3A 抗原（重组蛋白）

英文名称： FCGR3A Antigen (Recombinant Protein)

别名： Fc fragment of IgG, low affinity IIIa, receptor (CD16a); CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRIII; FCGRIII; FCRIIIA

储存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to a region derived from 17-208 amino acids of human FCGR3A

技术规格

Full name:	Fc fragment of IgG, low affinity IIIa, receptor (CD16a)
Synonyms:	CD16; FCG3; CD16A; FCGR3; IGFR3; IMD20; FCR-10; FCRIII; FCGRIII; FCRIIIA
Swissprot:	P08637
Gene Accession:	BC033678
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	This gene encodes a receptor for the Fc portion of immunoglobulin G, and it is involved in the removal of antigen-antibody complexes from the circulation, as well as other antibody-dependent responses. This gene (FCGR3A) is highly similar to another nearby gene (FCGR3B) located on chromosome 1. The receptor encoded by this gene is expressed on natural killer (NK) cells as an integral membrane glycoprotein anchored through a transmembrane peptide, whereas FCGR3B is expressed on polymorphonuclear neutrophils (PMN) where the receptor is anchored through a phosphatidylin

ositol (PI) linkage.