

兔抗 WDR77 多克隆抗体

中文名称: 兔抗 WDR77 多克隆抗体

英文名称: Anti-WDR77 rabbit polyclonal antibody

别名: p44; MEP50; MEP-50; HKMT1069; Nbla10071; p44/Mep50

储存: 冷冻 (-20℃)

抗原: WDR77

宿主: Rabbit

反应种属: Human, Mouse, Rat

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:	The protein encoded by this gene is an androgen receptor coactivator that forms a complex with protein arginine methyltransferase 5, which modifies specific arginines to dimethylarginines in several spliceosomal Sm proteins. The encoded protein may be involved in the early stages of prostate cancer, with most of the protein being nuclear-localized in benign cells but cytoplasmic in cancer cells. Several transcript variants encoding different isoforms have been found for this gene.
Applications:	ELISA, WB, IHC
Name of antibody:	WDR77
Immunogen:	Fusion protein of human WDR77
Full name:	WD repeat domain 77

Synonyms:	p44; MEP50; MEP-50; HKMT1069; Nbla10071; p44/Mep50
SwissProt:	Q9BQA1
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	100-300
WB Predicted band size:	37 kDa
WB Positive control:	LNCAP cell lysate
WB Recommended dilution:	500-2000



