

## CRYGS 抗原（重组蛋白）

中文名称： CRYGS 抗原（重组蛋白）

英文名称： CRYGS Antigen (Recombinant Protein)

别名： CRYG8; CTRCT20

储存： 冷冻（-20℃）

相关类别： 抗原

### 概述

Full length fusion protein

### 技术规格

<b>Full name:</b>	crystallin gamma S
<b>Synonyms:</b>	CRYG8; CTRCT20
<b>Swissprot:</b>	P22914
<b>Gene Accession:</b>	BC069478
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	Crystallins are separated into two classes: taxon-specific, or enzyme, and ubiquitous. The latter class constitutes the major proteins of vertebrate eye lens and maintains the transparency and refractive index of the lens. Since lens central fiber cells lose their nuclei during development, these crystallins are made and then retained throughout life, making them extremely stable proteins. Mammalian lens crystallins are divided into alpha, beta, and gamma families; beta and gamma crystallins are also considered as a superfamily. Alpha and beta families are f

urther divided into acidic and basic groups. Seven protein regions exist in crystallins: four homologous motifs, a connecting peptide, and N- and C-terminal extensions. Gamma-crystallins are a homogeneous group of highly symmetrical, monomeric proteins typically lacking connecting peptides and terminal extensions. They are differentially regulated after early development. This gene encodes a protein initially considered to be a beta-crystallin but the encoded protein is monomeric and has greater sequence similarity to other gamma-crystallins. This gene encodes the most significant gamma-crystallin in a adult eye lens tissue. Whether due to aging or mutations in specific genes, gamma-crystallins have been involved in cataract formation.