

C7 抗原(重组蛋白)

中文名称: C7 抗原(重组蛋白)

英文名称: C7 Antigen (Recombinant Protein)

别 名: complement component 7

储 存: 冷冻(-20℃)

相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 644-843 amino acids of human ${\rm C7}$

技术规格

Full name:	complement component 7
Swissprot:	P10643
Gene Accession:	BC063851
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	The complement cascade is a multi-protein system that functions to clear pathogens from an infected host. Part of the innate (unchangin g) immune system, the complement cascade consists of proteins and inactive zymogens that are present in blood and are stimulated by o ne of several triggers. Once stimulated, the cascade relays amplified r esponses throughout the body, ultimately activating the cell-killing m embrane attack complex which can insert itself into the cell membran e and cause the cell to lyse. C7 (complement component 7) is an 84 3 amino acid secreted protein that participates in the formation of m embrane attack complex (MAC), a complex that forms pores in the pl asma membrane of target cells for innate and adaptive immune resp



www.shjning.com

onses. As a membrane anchor, C7 exists as a monomer or dimer and can form multimeric rosettes with C5β. C7 defects are the cause of conent C7 deficiency (C7D), characterized by recurrent bacterial infections caused by Neisseria meningitidis.