

## CCDC112 抗原(重组蛋白)

中文名称: CCDC112 抗原(重组蛋白)

英文名称: CCDC112 Antigen (Recombinant Protein)

别 名: coiled-coil domain containing 112; MBC1

储 存: 冷冻 (-20℃)

相关类别: 抗原

概述

Fusion protein corresponding to a region derived from 247-446 amino acids of human CCDC112

## 技术规格

| Full name:         | coiled-coil domain containing 112   |
|--------------------|---|
| Synonyms:          | MBC1  |
| Swissprot:         | Q8NEF3  |
| Gene Accession:    | BC031242  |
| Purity:            | >85%, as determined by Coomassie blue stained SDS-PAGE  |
| Expression system: | Escherichia coli  |
| Tags:              | His tag C-Terminus, GST tag N-Terminus  |
| Background:        | CCDC112 (coiled-coil domain containing 112), also known as MBC1 (mutated in bladder cancer 1), is a 446 amino acid protein. The ge ne encoding CCDC112 is located on chromosome 5. Due to alterna tive splicing events, CCDC112 exists as two isoforms. Chromosome 5 comprises about 6% of human genomic DNA and contains 181 million base pairs encoding around 1,000 genes. It is associated wi th Cockayne syndrome through the ERCC8 gene and familial adeno matous polyposis through the adenomatous polyposis coli (APC) tu mor suppressor gene. Treacher Collins syndrome is also chromoso |



## www.shjning.com

me 5 associated and is caused by insertions or deletions within th e TCOF1 gene. Deletion of the p arm of chromosome 5 leads to C ri du chat syndrome. Deletion of 5q or chromosome 5 altogether i s common in therapy-related acute myelogenous leukemias and m yelodysplastic syndrome.